

Rassen, die eventuell vor Pupillendilatation untersucht werden sollten:

1. Iridocorneal angle abnormality using gonioscopy:

American Cocker Spaniel  
Bouvier des Flandres Bassets (all)  
Bloodhound Border Collie  
Chow Chow  
Dandie Dinmont Terrier  
Dutch Shepherd (Rough Hair)  
English Springer Spaniel  
Entlebucher Mountain Dog  
Flat Coated Retriever  
Golden Retriever  
Siberian Husky  
Leonberger  
Magyar Vizsla  
Samoyed Tatra

2. Persistent Pupillary Membrane using slitlamp biomicroscopy

Basenji  
Chow Chow  
English Cocker Spaniel  
Petit Basset Griffon Vendéen

3. Iris hypoplasia using slitlamp biomicroscopy

Australian Shepherd  
Dalmatian Rottweiler

4. Lens luxation (PLL)/KCS/vitreous degeneration/ocular melanosis using slitlamp biomicroscopy

Chinese Crested dog (PLL, vitreous degeneration, KCS)  
English Bulldog (KCS)  
Lancashire Heeler (PLL)  
Longhaired Dachshund (KCS)  
Lhasa Apso (KCS) Pug (KCS)  
Shih Tzu (KCS)  
Cairn Terrier also (ocular melanosis)  
Small Terrier breeds (PLL)  
West Highland White Terrier (KCS)  
Other breeds listed in Chapter 10 for PLL